

Assessment the Effect of *Spirulina platensis* as Supplemental Feed on Growth Performance and Survival Rate in Angel Fish (*Pterophyllum scalare*)

BibiMahsa Javadi Moosavi and Salar Montajami

Department of Fishery,

Gorgan University of Agricultural Sciences and Natural Resources, Gorgan, Iran

Abstract: This study was carried out to determine the effects of *Spirulina platensis* on the growth and survival rate in Angel fish (*Pterophyllum scalare*) as a protein source in a 2 months feeding trial. *S. platensis* was added to the basal diet at 0% (control), 5, 1 and 20% and fed to Angel fish (*Pterophyllum scalare*). The growth of Angel fish (*Pterophyllum scalare*) was related to the level of *S. platensis* containing. The final weight gain, specific growth rate, feed conversion ratio of fish were affected by spirulina supplementation ($p>0.05$). The highest Feed Conversion Ratio (FCR), 1.24 ± 0.79 was found in the fishes that were cultured with 5% of *S. platensis*. The survival rate was not affected by different amount of spirulina. The results of the present study, clearly show that *S. platensis* could be considered as a protein source for incorporation in Angel fish (*Pterophyllum scalare*) diets to gain more weight.

Key words: *Spirulina platensis*, Angel fish (*Pterophyllum scalare*), growth performance, survival rate, source, feed conversion ratio,

INTRODUCTION

Ornamental fish farming is an important primary industry. Ornamental fishes are often referred as living jewels due to their color, shape and behavior. They are peaceful, generally tiny, attractively colored and could be accommodated in confined space (Kasiri *et al.*, 2011; Lim and Wong, 1997). Modern ornamental fish culture and breeding operations have become vertically and horizontally intensified, necessitating a continuous supply of nutritionally balanced, cost effective feed (Kasiri *et al.*, 2011; Mandal *et al.*, 2010).

Microalgae are the natural foods of all aquatic animals. The use of microalgae in aquaculture is favored by several factors: Microalgae are the natural feeds of many aquaculture species and the basis of the natural food chains on which such species depend in the wild. Microalgae have high nutrient value, containing specialty feed components including pigments essential fatty acids and vitamins (Benemann, 1992). The use of microalgae in aquaculture has potential advantages due to the high conversion efficiencies (Benemann, 1992). Microalgae and their products support fish growth and it might be practical to feed microalgae to fish with little or no processing (Stanley and Jones, 1976).

Spirulina is a source of phycobiliproteins that are used as fluorescent markers in biomedical research. Earlier

studies examined how supplementing the diet with dry Spirulina powder affected the taste and quality of fishes (El-Sayed, 1994; Mustafa *et al.*, 1994; Watanabe *et al.*, 1990). Watanabe *et al.* (1990, 1993) reported that 5% dietary Spirulina supplementation depressed the lipid in the muscle and improved the taste and texture of striped jack *Pseudocaranx dentex*. Other studies suggested that using Spirulina in the feed of sea breams resulted in a high value product with a good market and high prices, apparently because the Spirulina significantly increased the stromal fraction which mainly contains collagen (El-Sayed, 1994; Mustafa *et al.*, 1994). In the case of ayu *Plecoglossus altivelis*, it has been claimed that fish grown on feed containing Spirulina are of better quality with better flavor, firmer flesh and brighter skin color (Mori *et al.*, 1987).

As the pigmentation additive, Spirulina was also found to improve the color of Angel fish. The feeding raw Spirulina as uni-feed to *Pterophyllum scalare* resulted in slightly better evaluations of color, texture and fatness than were obtained with commercial diets. The taste and smell evaluations were not different between the fish fed the Spirulina and those fed commercial diets (Lu and Takeuchi, 2002; Lu *et al.*, 2003). Only few data on growth performance are available for Angel fish fed with diet containing *S. platensis*.

The objective of this present investigation was study the efficacy of a fresh cell of cyanobacterium (blue-green

algae), *Spirulina platensis* as a feed ingredient on growth and survival of Angel fish in order to evaluate the suitability of the cyanobacterium as a feed ingredient.

MATERIALS AND METHODS

Experimental unit: About 10 aquaria of 85×45×55 cm (length×width×depth) was prepared at a private fishery laboratory in Mashhad, Iran.

Experimental animal: The fish were received from a public fish market in Mashhad. A total of 240 fish, the average weight of each fish before the experiment was 2.2±0.30 g and total length of 3.7±0.4 cm.

Experimental diets: The different feeding combinations (4 formulas of diets) were prepared as follows:

- Diet 1; the combination of feed containing 100% fish meal (T₁ or control)
- Diet 2; the combination of feed-stuff supplemented with 5% dried Spirulina powder (T₂)
- Diet 3; the combination of feed-stuff supplemented with 10% dried Spirulina powder (T₃)
- Diet 4; the combination of feed with 25% dried Spirulina powder (T₄)

Preparation of experimental diet: Well mixed feeding materials were packed in plastic bags and kept in the refrigerator at -18°C throughout the experiment. Micro-Kjeldahl methods were used to analyze protein content in feeds (AOAC, 1975). Fish were fed two times each day at 3% of Body Weight per day (BW/day) and the feeding rates were adjusted fortnightly.

Feed analysis: Nutrient compositions of experimental diets (Biomar) are given in Table 1 and typical nutritional composition of spray-dried spirulina powder is in Table 2. Proximate composition of diets was carried out using the association of analytical chemists (AOAC, 1975) methods. Protein was determined by measuring nitrogen (N×6.25) using the Kjeldahl method, crude fat was determined using petroleum ether (40-60 Bp) extraction method with Soxhlet apparatus and ash by combustion at 550°C.

Determination of growth parameters: Growth parameters were calculated as follows (Tacon, 1990; Hevroy *et al.*, 2005; Ai *et al.*, 2006):

$$\text{Body Weight Gain (BWG)} = \frac{\text{Final fish weight (G)} - \text{Initial fish weight (G)}}{\text{Initial fish weight (G)}}$$

Table 1: Nutrient composition of experimental diets (%)

Ingredients	Percentage
Protein	54.0
Lipid	18.0
Fiber	1.5
Ash	10.0
Vitamin	2.0

Table 2: Typical nutritional composition of spray-dried spirulina powder

Ingredients	Percentage
Crude protein	60.0
Carbohydrate	20.0
Fats	5.0
Minerals	9.0
Moisture	6.0
Carotenoids (red/yellow)	0.3
Chlorophyll (green)	1.0
Phycocyanin (blue)	12.5

$$\text{Specific Growth Rate (SGR)} = \frac{(\ln W_t - \ln W_0) \times 100}{t - 1}$$

$$\text{Survival rate} = \left(\frac{N_t}{N_0} \times 100 \right)^{\frac{1}{t}}$$

Statistical analysis: In order to determine significant differences, results were analyzed by one-way Analysis of Variance (ANOVA) and Duncan's multiple range tests were used to analyze the significance of the difference among the means of treatments by using the SPSS-17 programmer.

RESULTS AND DISCUSSION

Growth factors are summarized in Table 3. Results clearly show that the feeding rate showed a remarkable increment at 20% of *S. platensis* over the control. It, however followed a decreasing trend beyond 20% of Spirulina.

Fish larvae can be fed Spirulina either through gut-loaded rotifers and Artemia or through formulated microparticulate diets. Spirulina powder mixes easily with other ingredients for pelletizing. Care should be taken not to overheat the product. Temperatures high enough to denature carbohydrates will destroy the beneficial properties found in Spirulina. The basic method may be adapted to other species and local conditions. Best results are achieved when Spirulina is used in combination with fat-rich, live algae like diatoms. Spirulina powder is very rich in protein and vitamins but does not contain enough fat for growing larvae. Fat-rich diatoms, together with dried Spirulina can form an ideal diet that results in very strong postlarvae. Dried Spirulina may be used at each feeding or the hatchery operator may want to alternate it with a high quality, artificial microparticulate diet and Artemia, depending on local conditions and availability.

Table 3: Growth parameters and survival rate in Angel fish (*Pterophyllum scalare*) in the experimental treatments

Parameters	Control	5 (%)	10 (%)	20 (%)
Initial weight (g)	2.2±0.30	2.2±0.30	2.2±0.30	2.2±0.30
Final body weight (g)	3.9±0.04 ^c	4.8±0.6 ^a	4.7±0.03 ^a	4.2±0.07 ^b
Body weight gain (g)	1.70±0.26 ^c	2.60±0.3 ^a	2.5±0.27 ^a	2.0±0.23 ^b
Specific growth rate for weight (% BW day ⁻¹)	0.92±0.2 ^c	1.24±0.79 ^a	1.21±0.41 ^a	1.03±0.10 ^b
Survival rate (%)	100 ^a	100 ^a	100 ^a	100 ^a

Data are represented as Mean±SD; Means with the same letters in the same row are not significantly different

The present study indicates that the increased feed intake coupled with the improved food conversion could have enhanced the growth rate in the young ones of *Pterophyllum scalare* while it is very well clear that the Spirulina treated fish have better growth rate than the control, the maximum growth, however is achieved from 20%. Next to 20 and 10% was found to be effective. However, the under 5% showed a declining trend in growth parameters.

Cyanobacteria have been found to be a good source of protein for fish (Nandesha *et al.*, 2001). In addition, cyanobacteria have been reported to have no cell wall which results in improved digestion and absorption (Desikachary, 1959). The growth of Angel fish being positively affected at all levels of *S. platensis* inclusion. The difference in the FCR may also be due to a difference in the chemical composition of the diets.

No significant different of survival of Angel fish in this study possibly due to high protein content (54%) of the control diet that was enough to enhance immunity of the fish. The growth and survival rate of Angel fish being positively affected at all levels of *S. platensis* inclusion, thus clearly indicate the suitability of *S. platensis* for incorporation in the diet.

CONCLUSION

Spirulina is a rich source of protein, vitamins, minerals and pigments (Table 2). It also enhances the non-specific immune system, increasing the state of readiness of natural defenses. The combination of nutrients, pigments and immunostimulants explains reports of dietary Spirulina yielding better growth, reduction of stress and better appearance. Angel fish fed diet containing 5% of *S. platensis* gave the best result of growth, FCR. The results of the study clearly demonstrate that few cyanobacterium *S. platensis* could be used as a protein and lipid sources to incorporate into Angel fish diets to have a better culturing and gain market acceptance.

REFERENCES

- AOAC, 1975. Official Methods of Analysis. Association of Official Analytical Chemists, Washington DC., USA.
- Ai, Q., K. Mai, B. Tan, W. Xu, Q. Duan, H. Ma and L. Zhang, 2006. Replacement of fish meal by meat and bone meal in diets for large yellow croaker, *Pseudosciaena crocea*. Aquaculture, 260: 255-263.
- Benemann, J.R., 1992. Microalgae aquaculture feeds. J. Applied Phycol., 4: 233-245.
- Desikachary, T.V., 1959. Cyanophyta. 1st Edn., Indian Council of Agriculture Research, New Delhi, India, Pages: 686.
- El-Sayed, M.A.F., 1994. Evaluation of soybean meal, Spirulina meal and chicken offal meal as protein sources for silver seabream (*Rhabdosargus sarba*) fingerlings. Aquaculture, 127: 169-176.
- Høyvold, E.M., M. Espe, R. Waagbø, K. Sandnes, M. Ruud and G.I. Hemre, 2005. Nutrient utilization in Atlantic salmon (*Salmo salar* L.) fed increased levels of fish protein hydrolysate during a period of fast growth. Aquacult. Nutr., 11: 301-313.
- Kasiri, M., A. Farahi and M. Sudagar, 2011. Effects of feeding frequency on growth performance and survival rate of angel fish, *Pterophyllum scalare* (Perciformes: Cichlidae). Vet. Res. Forum, 2: 97-102.
- Lim, L.C. and C.C. Wong, 1997. Use of the Rotifer, *Brachionus calyciflorus* Pallas, in Freshwater Ornamental Fish Larviculture. In: Live Food in Aquaculture, Hagiwara, A., T.W. Snell, E. Lubzens and C.S. Tamaru (Eds.). Springer, Netherlands, ISBN: 9789048150038, pp: 269-273.
- Lu, J. and T. Takeuchi, 2002. Taste of tilapia *Pterophyllum scalare* fed solely on raw spirulina. Fish. Sci., 68: 987-988.
- Lu, J., T. Takeuchi and H. Ogawa, 2003. Flesh quality of tilapia *Pterophyllum scalare* fed solely on raw spirulina. Fish. Sci., 69: 529-534.
- Mandal, B., A. Mukherjee and S. Banerjee, 2010. Growth and pigmentation development efficiencies in fantail guppy, *Poecilia reticulata* fed with commercially available feeds. Agric. Biol. J. N. Am., 1: 1264-1267.
- Mori, T., T. Muranaka, W. Miki, K. Yamaguchi, S. Konosu and T. Watanabe, 1987. Pigmentation of cultured sweet smelt fed diets supplemented with a blue-green alga *Spirulina maxima*. Nippon Suisan Gakkaishi, 53: 433-438.

- Mustafa, M.G., T. Umino and H. Nakagawa, 1994. The effect of *Spirulina* feeding on muscle protein deposition in red sea bream, *Pagrus major*. *J. Applied Ichthyol.*, 10: 141-145.
- Nandesha, M.C., B. Gangadhara, J.K. Manissery and L.V. Venkataraman, 2001. Growth performance of two Indian major carps, catla (*Catla catla*) and rohu (*Labeo rohita*) fed on diets containing different levels of *Spirulina platensis*. *Bioresour. Tech.*, 80: 117-120.
- Stanley, J.G. and J.B. Jones, 1976. Feeding algae to fish. *Aquaculture*, 7: 219-223.
- Tacon, A.G.J., 1990. *Standard Methods for the Nutrition and Feeding of Farmed Fish and Shrimp*. Argent Laboratories Press, Washington, DC., USA., Pages: 454.
- Watanabe, T., W. Liao, T. Takeuchi and Y.H. Yamamoto, 1990. Effect of dietary *Spirulina* supplement on growth performance and flesh lipids of cultured striped jack. *J. Tokyo Univ. Fish.*, 77: 231-239.
- Watanabe, T., T. Takeuchi, N. Okamoto, S. Satoh, N. Iso, H. Sakamoto and T. Arakawa, 1993. Improvement of flesh quality of cultured striped jack with a newly developed soft-dry pellet. *J. Tokyo Univ. Fish.*, 80: 19-29.