

## Changes in Nutrient Composition, Trypsin Inhibitor, Phytate, Tannins and Protein Digestibility of Dolichos Lablab Seeds [Lablab Putrpuresus (L) Sweet] Occurring During Germination

Magdi A. Osman

Department of Food Science, College of Food and Agricultural Sciences,  
King Saud University, P.O. Box 2460, Riyadh 11451, Saudi Arabia

**Abstract:** The changes in nutrient composition, trypsin inhibitors, phytic acid, tannins and in-vitro digestibility of protein of lablab bean during 5 days of germination were studied. The crude protein contents was significantly increased, whereas lipid and carbohydrate contents decreased. Antinutritional study revealed that trypsin inhibitor activity and phytic acid content decreased as germination period increased. In contrast, there was a progressive increase in tannins content with increase in germination time. In-vitro protein digestibility markedly increased with germination time, with a significant increase in day 5.

**Key words:** Dolichos lablab, trypsin inhibitor, phytic acid, tannins, *in vitro* protein digestibility

### INTRODUCTION

Plants are important sources of energy and dietary proteins and play a significant role in human nutrition, particularly in the developing world. Among the plant sources, dry legumes and their products are among the richest sources of food protein. They are generally well adapted to a wide range of climates and environmental conditions. Dolichos lablab bean is an indigenous legume of the arid and semi-arid area of the South and East Asia where, it is traditionally consumed by the natives. Lablab bean plant produced well with limited moisture and is tolerant to temperature stress. It is regarded as potential food crop for arid lands of Africa, Middle East and South America. Lablab bean contains 22.4-31.3% protein and about 46.0-63.3%, 55% carbohydrates (Al-Othman, 1999; Deka and Sarkar, 1990; Ahamad and Nour, 1990; Salimath and Tharanathan, 1982). The bean can be consumed fresh, germinated, boiled steamed or fried. As in other legumes, Dolichos lablab bean contains some antinutritional components such as trypsin inhibitors, phytic acid and tannins (Vijayakumaris *et al.*, 1995; Devarj and Manjunath, 1995; Shastri and John, 1991). The nutritional quality and digestibility of plant protein is affected by presence of these antinutritional factors which may limit the use of plants as a source of protein. Several methods have been employed to improve the nutritional quality of legumes. The processing technologies employed include, soaking (Desphande and Sheryan, 1983; Vijayakumari *et al.*, 2007; Alonso *et al.*, 1998), heat treatment (Osman *et al.*, 2002; Di Pitro and Liener, 1989; Kadam and Smithard, 1987), fermentation (Reddy and Salunke, 1980; Osman, 2004) and irradiation (Abu Tarboush, 1998; Sidduraju *et al.*, 2002).

In the Orient and Far East Asia, germination process is used widely and routinely used as treatment to grain legumes. Germination, of the dry bean is reported to decrease the level of trypsin inhibitors (Frias *et al.*, 1995; Sathe *et al.*, 1983; Alonso *et al.*, 2000), phytic acid (Kataria *et al.*, 1989; Vidal-Valverde *et al.*, 1994; Sangronis and Machado, 2007) and tannins (Reddy *et al.*, 1985; El-Awady, 2002; Mubarak, 2005). Germination is also thought to increase *in-vitro* protein digestibility (Kataria *et al.*, 1989; Chau and Cheung, 1997) and essential nutrients (Ghorpade and Kadam, 1989). Contradictory reports on antinutritional factors in germinated seed are also available. Shastri and John (1991) observed an increase in trypsin inhibitors, phytic acid and tannins level during germination of Dolichos lablab. The sprouting, of sorghum, also produced an increase in tannins content (Ahamad *et al.*, 1996; Al-Jasser, 2005). This controversy as to whether the germination increase or decreases antinutritional factors seemed arisen because of different types of legumes or different germination conditions used in the studies. The objective of this study was to have a better understanding on the effect of germination on nutritive value, antinutritional factor level and in-vitro protein digestibility of Dolichos lablab.

### MATERIALS AND METHODS

**Materials:** Dolichos Lablab beans [*Lablab purpureus* cl.) sweet] were obtained from The Agricultural Experiment Station, College of Agriculture, King Saud University, Saudi Arabia. The Beans were cleaned manually and ground in a hammer mill to pass through No 30. Sieve.

**Germination:** The germinated beans were obtained by soaking the beans in water for 20 h. The soaked beans were spread on metal box, the lower side of which was a mesh covered with a piece of jute sack, another piece of jute sack was used to cover the box. The box was kept at room temperature in the dark. The beans were watered once a day. Germination was carried out for 5 days.

**Sampling, drying and grinding:** Samples of germinated beans were taken daily for subsequent analysis. Drying of the germinated beans was carried out at room temp. The dried germinated beans, were ground using a hammer mill, to pass through 30 mesh screen size.

**Proximate composition:** The protein, fat, ash, crude fiber and moisture content of the bean samples were estimated according to the method described by the AOAC (1984). Total carbohydrates were calculated by difference.

**Trypsin Inhibitor Activity (TIA):** Trypsin inhibitor activity was assayed according to Kakade *et al.* (1969) using BAPA N-benzoyl-DL-arginine-P-nitroanilide hydrochloride and trypsin type III from bovine pancreas. TIA expressed as trypsin inhibitor unit per milligram of sample (TIU mg<sup>-1</sup> sample) was calculated from the absorbance reading against a blank in spectrophotometer. One trypsin unit is defined as an increase of 0.01 in absorbance reading at 410 nm per 10 mL of the reaction mixture.

**Phytic acid:** Phytic acid analysis was performed according to the method of Latta and Eskin (1980). Duplicate fat free dry samples weighing 5 g were extracted in 100 mL of 0.667 N HCL for 1 h at room temp. The extract was filtered under vacuum. The phytate content in sample extract was determined by ion exchange method. Phytic acid in elute was determined by a colorimetric method.

**Tannin:** The modified vanillin-HCl method of Price *et al.* (1978) was followed with minor modification. One gram of sample was extracted with 10 mL 1% HCl in methanol for 24 h at room temperature then centrifuged at 5000 rpm. Vanillin HCl reagent was prepared by mixing prior to use equal volumes of 8% HCl in methanol with 2% vanillin in methanol. One milliliter of supernatant was mixed with 5 mL of vanillin HCl reagent. The absorbance was read at 500 nm after 20 min incubation at room temperature.

**In vitro protein digestibility:** *In vitro* digestibility was determined following Hsu *et al.* (1977) as modified by Satterlee *et al.* (1979). The drop of pH of casien (control) and the sample after 20 min hydrolysis by proteolytic

enzymes was measured using an Orion pH meter. The enzymes used were trypsin type IX from porcine pancreas, chymotrypsin type II from bovine pancreas, peptidase type III from porcine intestine and protease type VI from streptomyces griseus. All enzymes were supplied by Sigma chemical company (St. Louis, Mo. USA). The *in vitro* digestibility was calculated according to Satterlee *et al.* (1979) equation.

$$\% \text{ In-vitro digestibility} = 234.84 - 22.56 X$$

Where, X = the pH of suspension after 20 min hydrolysis of protein.

**Statistical analysis:** The data were analyzed used one way ANOVA with mean separated by Least Significance Differences (LSD) at  $p < 0.05$  (Steel and Torrie, 1980).

## RESULTS AND DISCUSSION

**Proximate composition:** Table 1 Shows the changes in proximate composition during the germination period of lablab beans. There was a gradual increase in moisture with the time of germination, with 5th day showing the highest moisture content. The moisture content in lablab bean increased from 6.41-12.93% during germination period. The increase in moisture content is related to increase in water activity during germination due to hydrolytic enzymes. Protein content of germinated bean showed significant decrease in the beginning and then gradual increase was observed. At the end of germination protein content significantly increased. This increase, attributed to the utilization of carbohydrates as source of energy during germination process. The results of this work were in agreement with those obtained with winged bean (King and Putwastien, 1987), Nigerian cowpea (Akpapunam and Achinewhu, 1985), Tepary bean (Idouraine *et al.*, 1989), faba and kidney bean (Alonso *et al.*, 2000) and mung bean (Mubarak, 2005). Fat content of lablab bean decreased significantly by approximately 37% following germination. The reduction was probably caused by break-down of fat by beta oxidation with fat being used for energy purposes in embryo development. These observations, are in agreement with those reported by Idouraine *et al.* (1989), Alonso *et al.* (1998), El-Awady (2002) and Mubarak (2005) and Ghavidel and Prakash (2007). Crude ash content did not vary significantly between the raw and germinated samples. There was a significant increase in total carbohydrates in lablab bean germinated for 48 h, then a decrease occurred after this period. The reduction in total carbohydrates was attributed to the increase in amylotic

**Table 1: Proximate composition of raw and germinated seed of the lablab bean on dry weight basis**

	Moisture	Protein	Fat	Ash	Carbohydrates
Raw	6.41±0.03 <sup>D</sup>	26.86±0.3 <sup>B</sup>	1.9±0.01 <sup>A</sup>	3.96±0.04 <sup>BC</sup>	67.23±0.30 <sup>CD</sup>
First day	7.79±0.07 <sup>BC</sup>	25.80±0.13 <sup>DC</sup>	0.87±0.13 <sup>D</sup>	3.46±0.04 <sup>BC</sup>	69.82±0.23 <sup>A</sup>
Second day	7.93±0.02 <sup>B</sup>	25.07±0.97 <sup>DEF</sup>	0.86±0.04 <sup>D</sup>	3.70±0.01 <sup>BCD</sup>	70.38±1.00 <sup>A</sup>
Third day	7.80±0.08 <sup>BC</sup>	26.60±0.32 <sup>BC</sup>	0.86±0.13 <sup>D</sup>	3.69±0.02 <sup>BCD</sup>	68.46±0.42 <sup>B</sup>
Fourth day	7.60±0.80 <sup>BC</sup>	27.08±0.47 <sup>B</sup>	0.97±0.03 <sup>DC</sup>	3.66±0.01 <sup>BCD</sup>	68.27±0.42 <sup>CD</sup>
Fifth day	12.95±0.80 <sup>A</sup>	28.55±0.21 <sup>A</sup>	1.19±0.08 <sup>BC</sup>	3.83±0.03 <sup>ABC</sup>	66.40±0.14 <sup>D</sup>

All values are means of 3 replicates±SD, Values with same letters (a, b, c, d, e, f within columns) are not significantly different at  $p < 0.05$

**Table 2: Change in antinutritional factors level and protein digestibility during germination of the lablab bean**

	Trypsin inhibitor (TIU mg <sup>-1</sup> )	Phytic acid (mg 100 gm <sup>-1</sup> )	Tannins (%catechin equivalent)	IVPD
Raw	28.96±0.30 <sup>A</sup>	605.39±0.39 <sup>A</sup>	0.42±0.01 <sup>F</sup>	88.17±1.70 <sup>BC</sup>
First day	26.85±0.23 <sup>B</sup>	406.69±1.66 <sup>B</sup>	0.56±0.02 <sup>DE</sup>	88.27±0.00 <sup>BC</sup>
Second day	25.07±0.30 <sup>C</sup>	393.24±3.84 <sup>C</sup>	0.57±0.01 <sup>BDE</sup>	88.43±0.00 <sup>BC</sup>
Third day	24.61±0.31 <sup>D</sup>	358.87±4.06 <sup>D</sup>	0.60±0.014 <sup>C</sup>	90.36±0.56 <sup>AB</sup>
Fourth day	23.51±0.99 <sup>E</sup>	317.72±5.01 <sup>E</sup>	1.15±0.01 <sup>B</sup>	89.33±1.73 <sup>AC</sup>
Fifth day	23.35±0.81 <sup>E</sup>	309.10±5.35 <sup>E</sup>	1.27±0.01 <sup>A</sup>	92.27±1.83 <sup>A</sup>

All values are means of 3 replicates±SD, Values with same letters (a, b, c, d, e, f within columns) are not significantly different at  $p < 0.05$

enzyme activity that hydrolyzes starch into simple absorbable sugar to the developing embryo. Similar results were reported by Ologhobo and Fetuga (1986) for germinated cowpea, Mbithi-Mwikya *et al.* (2000) for germinated finger millet, El-Adway (2002) for germinated chickpea and Mubarak (2005) for germinated mung bean.

**Trypsin inhibitor:** The change in Trypsin Inhibitor Activity (TIA), Phytic Acid (PA) and tannins during germination is presented in Table 2. A significant ( $p < 0.05$ ) decrease in TIA level was observed at the 1st day of germination and further significant reduction following the increase of germination time. During 5 days germination period the TIA progressively decreased from 28.96-23.35 representing reduction of 19.4% at the end of the 5th day. The results obtained in this research agree with those reported for Kidney bean (El-Hag *et al.*, 1978), navy bean Great Northern bean (Sathe *et al.*, 1983), lima bean (Ologhobo and Fetuga, 1983; mung bean (Mubarak, 2005) lentil (EL-Mahdy *et al.*, 1983; Frias *et al.*, 1995; Valverde *et al.*, 1994), soybean (Collin and Sandars, 1976), pigeon bean (Torres *et al.*, 2007; Sangronis and Machado, 2007) and faba bean and kidney bean (Alonso *et al.*, 2000). Such pattern were also observed in other plant like sorghum (Al-Jasser, 2005), finger millet (Mbithi-Mwikya *et al.*, 2000), small radish, radish, white mustard and rapeseed (Frias *et al.*, 2005) and breadnut, cashew nut and fluted pumpkin (Fagbemi *et al.*, 2005). Our results with germinated lablab bean were not exactly the same as those reported earlier by Shastry and John (1991) which indicated TIA increase during the progressive of germination. Similarly, Chang and Herrold (1988) observed an increase in TIA in pinto bean during germination.

**Phytic acid:** There was a significant reduction in phytic acid content with the increase of germination time of the Dolichos lablab beans (Table 2). The percent of decrease

in phytic acid content after 5 days germination was 49.9%. This finding agree with that of Sathe *et al.* (1983), who reported 57.8 reduction in PA content of the Great Northern bean after 5 days of germination. Many workers noticed significant reduction in PA content of legumes. Alonso *et al.* (2000) reported 61 and 30% reduction in PA content in faba bean and kidney bean respectively after germination. Sangronis and Machado (2007) found that germination reduced PA content in white bean, black bean and pengion pea by 44.7, 52.9 and 40.7, respectively. Similarly, Ghavidel and Parkash (2007) observed 18, 20, 21.1 and 20.8% reduction PA 24 h germination for green gram, cowpea, lentil and chickpea, respectively. In contrast germination of Dolichos lablab for 8 days increased phytic acid content (Shastry and John, 1991). The phytate reduction caused by germination was probably due the phytase activity that hydrolyzed phytates.

**Tannins:** Data on the effect of germination on tannins content of Dolichos lablab, is shown in Table 2. In contrast to trypsin inhibitor and phytic acid, the tannins content significantly increased as period of germination increased. Similar observations were reported by many investigators. Shastry and John (1991) observed significant increase in tannin in Dolichos lablab after 8 day germination. Similarly Valverde *et al.* (1994) reported increase in tannins content in lentil after 6 days of germination. An increase in tannins content has also been reported for sorghum when sprouted for 5 days (Abawi *et al.*, 1986; Ahamad *et al.*, 1996; Al-Jaseer, 2005). However, several investigators have observed decrease in tannins content on various legumes due to germination. Reddy *et al.* (1985) found that germination for 2 days after soaking reduced the tannins content by 50% in a wide variety of legumes. Alonso *et al.* (2000) also reported a marked reduction in tannins after 72 h germination of

faba bean and kidney bean. Similarly, Shimelis and Rakshit (2006) reported reduction in tannins content in sprouted kidney bean. More recently Ghavidel and Praksh (2007) reported decrease in tannins content in green bean, cowpea, lentil and chickpea germinated for 24 h.

***In vitro* Protein Digestibility (IVPD):** There was a progressive increase in the protein digestibility as the period of germination increase (Table 2). The values for digestibility after 5 days were significantly ( $p>0.05$ ) higher than those after the first three days. These results indicate that the tannins content had no significant effect on *in-vitro* digestibility. The improvement in IVPD of *Dolichos lablab* during germination may be attributed to the modification and degradation of storage proteins. Germination has been reported to increase protein digestibility of *Dolichos lablab* (Shastri and John, 1991), of green gram, cowpea, lentil and chickpea (Ghavidel and Prakash, 2007), faba bean and kidney bean (Alonso *et al.*, 2000), white bean, black bean and pigeon bean (Sangronis and Machado, 2007), sorghum cultivars (ELmaki *et al.*, 1999; Al-Jasser, 2005), mungbean (Kataria *et al.*, 1989), Kidney bean (Shimelis and Rakshit, 2006).

## CONCLUSION

Germination resulted in increased level of the protein and reduced contents of the lipid and carbohydrates of *Dolichos lablab*. Germination was found to be effective in reducing trypsin inhibitor and phytic acid content. However, it was found to increase tannins content. Protein *in-vitro* digestibility values increased as germination period increase. Therefore, germination of *lablab* seed holds a good potential for improving nutritional value of the bean by reduction in antinutritional and increasing protein *in-vitro* digestibility and thereby enhancing its utilization.

## ACKNOWLEDGEMENT

The author is grateful to the Agriculture Research Center, College of Food Science and Agriculture, King Saud University for the financial assistance. The Author wish to thank Mr. Ibrahim El-Zien and Mr. Abubaker for their excellent technical assistanc.

## REFERENCES

Abawi, F.G., A.Y. Masaad and T.W. Sullivan, 1986. Nutritional value of pretreated sorghum in broiler diet. *Poult. Sci.*, 65: 1:1.

Abu-Tarboush, H.M., 1998. Irradiation inactivation of some antinutritional factors in plants seeds. *J. Agric. Food Chem.*, 46: 2966-2702.

Ahamed, S.B., A.S. Mahgoub and Babikir, 1996. Changes in tannins and cyanide contents and diastic activity during germination and the effect of traditional processing on cynide content of sorghum cultivars. *Food Chem.*, 56: 159-162.

Ahmed, A.H.R. and. A.A. Nour, 1990. Protein quality of common Sudanese leguminous seed. *Labensm Wiss U. Tech.*, 23: 301-304.

Akpapunam, M.A. and S.C. Achinewhu, 1985. Effect of cooking, germination and fermentation on chemical composition of Nigerian cowpea (*Vigna unguiculata*). *Qual. Plant. Foods Hum. Nutr.*, 35: 353-357.

Al-Jasser, S.M., 2005. Effect of soaking and germination time on antinutritional factors and protein digestibility of three sorghum varieties grown in Saudi Arabia. *Alexandria Sci. Exchange J.*, 26: 315-321.

Alonso, R., E.A. Aguirre and F. Marzo, 2000. Effect of extrusion and traditional processing methods, on antinutritional and *in vitro* digestibility of protein and starch in faba and kidney beans. *Food Chem.*, 68: 159-165.

Alonso, R., Orue and F. Marzo, 1998. Effect of extrusion and conventional processing methods on protein and antinutritioal factor content in pea seeds. *Food Chem.*, 63: 505-512.

Al-Othman, A.A., 1999. Chemical composition and nutritional evaluation of *Dilichos lablab* bean [*Lablab purpureus* (L.) sweet] grown Al-Gassim regionod Saudi Arabia. *Annala Agric. Sci., Ain Shims Univ. Cairo.*, 44: 641-652.

AOAC, 19884. Official Methods for Analysis (14th Edn.), Association of Official Analytical Chemists. Arlington VA. USA.

Chang, K.K. and R.L. Harrold, 1988. Change in selected biochemical components *in vitro* digestibility and amino acids in two bean cultivars during germination. *J. Food Sci.*, 53: 783-787.

Chau, C.R. and P.K. Cheung, 1997. Effect of various processing on antinutritional and *in vitro* digestibility of protein and starch of two Chinese indigenous legume seeds. *J. Agric. Food Chem.*, 45: 4773-4776.

Collins, J.L. and G.G. Sanders, 1976. Changes in TI activity in some soybean varities guring maturation and germination. *J. Food. Sci.*, 41: 167-172.

Deka, K.R. and C.R. Sarkar, 1990. Nutrient composition and antinutritional factors of *Dolichos lablab* L. seeds. *Food Chem.*, 38: 239-246.

- Desphande, S. 1983. Acid and tannins and trypsin inhibitor activity on soaked dry beans (*Phaseolus vulgaris* L.) *Nutr. Rep. Int.*, 27: 371-375.
- Devaraj, V.R. and N.H. Manjuntath, 1995. Effect of cooking on proteinase inhibitors of *Dolichos Lablab* bean (*Dolichos lablab prepureus*). *Plant. Food. Hum. Nutr.*, 48: 107-112.
- Di Pitero, C.M. and I.E. Liener, 1989. Heat inactivation of Kunitz and Bowman-Birk proteinase inhibitors. *J. Agric. Food Chem.*, 37: 39-44.
- El-Adawy, T.A., 2002. Nutritional composition and antinutritional factors of chickpea (*Cicer arietinum* L.) undergoing different cooking methods and germination. *Plant Food Human Nutr.*, 57: 83-97.
- El-Hag, N., N.F. Haard and R.E. Morse, 1978. Influence of sprouting on digestibility coefficient, trypsin inhibitor and globulin protein of red kidney bean. *J. Food Sci.*, 43: 1784-1880.
- EL-Mahdy, A.R., Y.G. Moharram and O.R. Abou-Samaha, 1983. Influence of germination on nutritional quality of lentil seeds. *Z. Lebensm. Unter. Forsch.*, 181: 318-320.
- Elmaki, B.H., E.E. Babiker and El Tinay, 1999. Change in chemical composition, grain malting, starch and tannin content and protein digestibility during germination of sorghum cultivars. *Food Chem.*, 64: 331-336.
- Fagbemi, T.N., A.A. Oshodi and K.O. Ipinmoroti, 2005. Processing effect on some antinutritional factors and *In Vitro* Multienzyme Protein Digestibility (IVPD) of three tropical seeds: Bread nut (*Artocarpus altilis*), cashewnut (*Anacardium occidentale*) and fluted pumpkin (*Telfairia occedentale*). *Pak. J. Nutr.*, 4: 250-256.
- Frias, J., H. Zielinski, M. Piskula, H. Kozłowska and C. Vidal-Valverde, 2005. Inositol phosphate content and trypsin inhibitor activity in ready to eat cruciferous sprouts. *Food Chem.*, 93: 331-336.
- Frias, J., C. Diaz-Pollan, C.L. Hedley and C. Vidal-Valverde, 1995. Evolution of trypsin inhibitor activity during germination of lentils. *J. Agric. Food Chem.*, 43: 2231-2234.
- Ghavidéh, R.A. and J. Prakash, 2007. The impact of germination and dehulling on nutrient, antinutrient *in vitro* iron and calcium bioavailability and *in vitro* starch and protein digestibility of some legume seeds. *LWT*, doi:10.1016/j.lwt.2005.08.002.
- Ghorpad, U.M. and S.S. Kadam, 1989. Germination. In *CRC. Handbook of World Legumes: Nutritional Chemistry, Processing Technology and Utilization*; Salunkhe, D.K., Kadam, S. S., (Eds.), CRC Press: Boca Raton, Fl., 3: 113-122.
- Hsu, H.W., D.L. Vavak, L.D. Satterlee and G.A. Miller, 1977. Multi-enzyme technique for estimating protein digestability. *J. Food Sci.*, 42: 1269-1273.
- Idouraine, A., A.M. Tinseley and C.W. Weber, 1989. Nutritional quality and sensory acceptability of akara prepared from germinated tepary bean. *J. Food Sci.*, 54: 114-117.
- Kadam, S.S. and R.R. Smithard, 1987. Effect of heat treatment on trypsin and hemagglutinating activities in winged bean. *Plant. Foods. Hum. Nutr.*, 37: 151-159.
- Kakade, M.L., N. Simon and I.E. Liener, 1969. An evaluation of natural vs. synthetic substrate for measuring antitryptic activity of soybean samples. *Cereal Chem.*, 46: 518-526.
- Kataria, A., B.M. Hauhan and D. Punia, 1989. Antinutrient and protein digestibility (*in vitro*) of mung bean as affected by domestic processing and cooking. *Food Chem.*, 32: 9-17.
- King, R.D. and P. Puwastien, 1987. Effect of germination on the proximate composition and nutrient quality of winged bean (*Psophocarpus tetragolobolus*) seeds. *J. Food. Sci.*, pp: 52-56.
- Latta, M. and M. Eskin, 1980. A colorimetric method for phytate determination. *J. Agric. Food Chem.*, 28: 1313-1315.
- Mbithi-Mwikya, S., J. van Camp, Y. Yiru and A. Huyghebaert, 2000. Nutrient and antinutrient changes, in finger millet (*Eleusine coracana*) during sprouting. *LWT.*, 33: 9-14.
- Mubarak, A.E., 2005. Nutritional composition and antinutritional factors of mung bean seeds (*Phaseolus aureus*) as affected by some home traditional processes. *Food Chem.*, 89: 489-495.
- Ologhobo, D.A. and B.L. Fetuga, 1983. Trypsin inhibitor activity in some lima bean varieties as affected, by different processing method. *Nutr. Rep. Int.*, 27: 41-48.
- Ologhobo, D.A. and B.L. Fetuga, 1986. Change in carbohydrate contents of germinating cowpea seeds. *Food Chem.*, 20: 117-120.
- Osman, A.M., 2004. Changes, in sorghum enzyme inhibitors, phytic acid, tannins and *in vitro* protein digestibility occurring during Khamir (local bread) fermentation. *Food Chem.*, 88: 129-134.
- Osman, A.M., M.R. Reid and C.W. Weber, 2002. Thermal inactivation of tepary bean (*Phaseolus acutifolius*), soybean and lima bean protease inhibitors: Effect of acidic and basic pH. *Food Chem.*, 78: 419-423.
- Price, M.L., V.C. Scoryoc and L.G. Butter, 1978. A citric evaluation of reaction as assay for tannin in sorghum grain. *J. Agric. Food Chem.*, 26: 1214-1218.

- Rahma, E.M., A.A. EL-Bawey, T.A. El-Adawy and M.A. Guma, 1987. Changes in chemical and antinutritional factors and functional properties of faba bean during germination. *Lebensm. Wiss. Tech.*, 20: 271-276.
- Reddy, N.R., Pierson, M.D., Sathe, S.K. and Salunkhe, D.K.. 1985. Dry bean tannin: A review of nutritional implication. *J. Am. Oil. Chem. Soc.*, 62: 451-549.
- Reddy, N.R. and D.K. Salunke, 1980. Effect of fermentation, on phytate phosphorus and mineral of black gram, rice and black gram-rice blend. *J. Food Sci.*, 45: 1708-1712.
- Salimath, V.P. and R.N. Tharanathan, 1982. Carbohydrates of field bean (*Dolichos lablab*). *Cereal Chem.*, 59: 430-435.
- Sangronis, E. and C.J. Machado, 2007. Influence of germination on the nutritional quality of *Phaseolus vulgaris* and *Cajanus cajan*. *LWT.*, 40: 116-120.
- Sathe, S.K., S.S. Desphande, N.R. Reddy, D.E. Goll and D.K. Salunkhe, 1983. Effect of germination on proteins, raffinose oligosaccharides and antinutritional factors in Great Northern beans (*Phaseolus vulgaris* L.). *J. Food Sci.*, 48: 1796-1800.
- Satterlee, L.D., J.G. Kendrick and G.A. Miller, 1979. Measuring protein quality. *J. Am. Oil Chem. Soc.*, 56: 103-108.
- Shastri, M. and E. John, 1991. Biochemical change and *in-vitro* protein digestibility of endosperm of germinating *Dolichos lablab*. *J. Sci. Food Agric.*, 55: 529-538.
- Shimelis, E.A. and S.K. Rakshit, 2006. Effect of processing on antinutritional and *in vitro* protein digestibility of kidney bean (*Phaseolus vulgaris* L.) varieties grown in East Africa. *Food Chem.*, doi:10.1016/J. Food Chem., 2006.08.005.
- Sidduraju, P., O. Osoniyi, H.P.S. Makkar and K. Becker, 2002. Effect of soaking and ionizing on various antinutritional factors of seeds from different sp. of an unconventional legumes, *Sesbana* and common legume, green gram (*Vigna radiate*). *Food. Chem.*, 79: 273-281.
- Steel, R.G. and J.H. Torri, 1980. Principles and procedures of statistics. Mc Graw- Hill. Book Co. Inc., New York, USA.
- Torres, A., J. Frias, M. Granito and C. Vidal-Valverde, 2007. Germinated *Cajanus cajan* seeds as ingredient in pasta products: Chemical and biological and sensory evaluation. *Food Chem.*, 101: 202-211.
- Vidal-Valverde, C., J. Frias, I. Estrella, J.M. Gorospe, R. Ruiz and J. Bacon, 1994. Effect of processing on some antinutritional factors of lentils. *J. Agric. Food. Chem.*, 42: 2291-2295.
- Vijayakumari, K., M. Pugalenth and V. Vadivel, 2007. Effect of soaking and hydrothermal processing methods on the level of antinutritional and *in vitro* protein digestibility of *baubinia purpurea* L. seeds. *Food Chem.*, 103: 968-975.
- Vijayakumari, K., P. Siddhuraju and K. Janardanan, 1995. Effect of various water or hydrothermal treatments on certain antinutritional compounds in the seed of tribal pulse, *Dolichos lablab* var. *vulgaris* L. *Plant. Food. Human Nutr.*, 48: 17-29.