

Composition and Abundance of Macroenthos in Majidun River, Ikorordu Lagos State, Nigeria

I.K. Esenowo and A.A.A. Ugwumba
Department of Zoology, University of Ibadan, Ibadan, Oyo State, Nigeria

Abstract: A study on the composition and abundance of macroenthos of Majidun river, Ikorodu, Lagos state was carried out from June, 2008 to May 2009 at 6 sampling stations along the river from its source to the mouth. The river is a multipurpose resource for artisanal fishing, transportation, sand mining and domestic uses. A total of 10,799 individuals of macroenthos belonging to 18 genera in two Phyla, Mollusca and Anthropoda were recorded from the 6 sampling stations. Mollusca were represented by Bivalvia and Gastropoda, the most common species were the gastropods: *Pachymelania aurita*, *P. fusca*, *P. fusca* var. *quadriserlata*, *Tympanotomus fuscatus* var. *radula*, *T. fuscatus fuscatus* and *Melanooides tuberculata*. Arthropoda consisted mainly of caddis fly larva, *Phryganea* sp., the crabs: *Sesarma huzardii*, *Cardisoma armatum* and *Callinectes latimanus* and the prawn: *Macrobrachium marobrachion*. *T. fuscatus* var. *radula* was the most abundant species accounting for 20.17%, closely followed by *P. aurita* 19.58% while the least abundant, *Phryganea* sp., accounted for <0.1% of the total number of macroenthos. Shannon-wiener diversity index indicated a maximum of 0.97 in mid river station (surrounded by mangrove trees) compared to the stations at the source of the river (0.88) and its mouth (0.87) where it flows into Lagos Lagoon. Macroenthos abundance and diversity levels were significantly low.

Key words: Macroenthos, composition, diversity, abundance, ikorodu, Majidun river

INTRODUCTION

Macroenthos play an important role in aquatic community which includes mineralization, mixing of sediments and flux of oxygen into sediment, cycling of organic matter and assessing the quality of inland water (George *et al.*, 2009). The distribution of macroenthos fauna are determined by a number of factors such as physical nature of the substratum, depth, nutritive content, degree of stability and oxygen content of the water body. Benthic microinvertebrates are threatened by changes in its habitat which are associated with pollution, erosion and siltation (Lydeard *et al.*, 2004).

In assessing the health of aquatic environment, bioassessment has become a reliable method for measuring human influence on aquatic ecosystems, complementing traditional physical and chemical methods. The presence or absence of benthic macroinvertebrates has been shown to be a good indicator of both chronic and episodic impact of human disturbance to river condition and other aquatic environment (Rosenberg and Resh, 1993).

Diaz *et al.* (2004) described benthic infauna as opportunistic species adapted to a dynamic salinity regime and variable physical conditions. Benthic species

composition, abundance, biomass and distribution patterns are to a large extent determined by a mixture of interacting variable of physico-chemical parameters like temperature regime, dissolved oxygen concentration, salinity and biochemical oxygen demand (Dillon, 2000; Edokpayi *et al.*, 2000; Celik, 2002; Egborge *et al.*, 2003; Ikomi *et al.*, 2005).

Majidun river is one of the important rivers in Lagos state found in the Western part of Nigeria. It is noted for artisanal fisheries and transportation. Macroenthos constitute the food for fishes hence, it is important to have reliable information on species composition. Several researches exist on the diversity and abundance of species composition for other freshwater system located in Lagos state such as the Lagos Lagoon (Akpata *et al.*, 1993; Brown and Ajao, 2004), Lekki Lagoon (Awosika and Dublin-Green, 1994) and Port Novo Creek (Chukwu and Nwankwo, 2003). This study constitutes the 1st record of an investigation into the diversity and abundance of macroenthos in Majidun river.

MATERIALS AND METHODS

Study area: Majidun river is located in Ikorodu, Lagos state on the Southwestern part of Nigeria coastline within




Fig. 1: Map of Ikorodu showing Majidun river and sampling stations (1-6)

longitude 3.22° and 3.29°E and latitude 6.39° and 6.41°N (Fig. 1). It has a catchment area of 2.9 km² with its major source from River Owuru and flows into Lagos Lagoon at Ipakodo from where it receives tidal waters from the Lagoon. The river is a multipurpose resource for artisanal fishing, sand mining, transportation and for domestic uses. Sandy beaches are seen in stations 4-6 of the river where there is sand mining activities while stations 1-3 are fringed with tidal mud flats and mangrove swamps in which plants such as *Rhizophora racemosa*, *Avicenia nitida* and *Echiorna crassepes* are to be found.

At each sampling station, three replicate samples of macrobenthos were collected monthly using 0.1 m² Van veen Grab from June 2008 to May 2009. The sediment collected were poured into polythene bags, labeled and brought to the laboratory for examination.

In the laboratory, each sediment sample was washed through three sets of sieves, 1st 2 mm then 1 mm and finally 0.5 mm mesh size, sieve to collect the macrobenthos in them. The retained macrobenthos were poured into a white enamel tray and stained with Rose Bengel solution (Holme and McIntyre, 1984). They were sorted using forceps, identified using Macan (1959), Pennak (1978), Edmunds (1978) and WHO (1978) and counted. Macrobenthos diversity was estimated using the Shannon-weiner index of diversity (H) (Shannon and Weaver, 1963) and Equitability index or Evenness (J) (Lloyd and Ghellardi, 1964).

RESULTS

The composition, abundance and distribution of the macrobenthos in the study area are shown in Table 1. A




Fig. 2: Percentage (number) of macrobrnths in Majidun river




Fig. 3: Monthly relative abundance of macrobenthos in Majidun river (July 2008 to June 2009)

total of 18 Macrobenthos genera belonging to two phyla Mollusca and Arthropoda were recorded. Phylum Mollusca was represented by gastropoda and bivalvia while Arthropoda were mainly decapods and trichoptera. The gastropoda, *Tympanotomus fuscatus* var. *radula* (L) was the most dominant species accounting for about

Table 1: Composition and relative abundance of macrobenthos at the 6 stations in Majidun river

Macrobenthos taxa	Station 1		Station 2		Station 3		Station 4		Station 5		Station 6	
	No	(%)	No	(%)	No	(%)	No	(%)	No	(%)	No	(%)
Gastropoda												
<i>Pachymelania aurita</i>	13	1.5	3	0.3	65	4.1	610	21.1	748	22.8	210	19.6
<i>Pachymelania fusca</i>	10	1.2	2	0.2	13	0.8	330	11.4	341	10.4	82	7.6
<i>Pachymelania fusca</i> var. <i>quadriseolata</i>	122	14.4	296	26.4	377	23.7	388	13.4	434	13.2	140	13.1
<i>Tympanotomus fuscatus</i> var. <i>radula</i> (L.)	281	33.1	251	22.4	377	23.7	547	19.0	459	14.0	264	24.6
<i>Tympanotomus fuscatus fuscatus</i>					8	0.5	112	3.9	148	4.5	49	4.6
<i>Melanoides tuberculata</i>	158	18.6	264	23.6	432	27.1	390	13.5	571	17.4	173	16.1
<i>Neritina afra</i>	30	3.5	32	2.9	49	3.1	64	2.2	99	3.0	50	4.7
<i>Lanistes varicus</i>	34	4.0	18	1.6	5	0.3	3	0.1	1	0.1	*	*
<i>Lanistes ovum</i>	14	1.6	5	0.4	4	0.3	4	0.1	1	0.1		
Bivalvia												
<i>Macra nitida</i>	17	2	17	1.5	21	1.3	55	1.9	49	1.5	16	1.5
<i>Diplodonta diaphana</i>	14	1.6	100	8.9	140	8.8	106	3.7	299	9.1	83	7.7
<i>Mytilus perna</i>	*	*	4	0.4	21	1.3	59	2.0	21	0.6		
<i>Arca senilis</i>	*	*	*	*	4	0.3	37	1.3	27	0.8	3	0.3
Decapoda												
<i>Sesarma angolense</i>	90	10.6	60	5.4	15	0.9	75	2.6	66	2.0	2	0.2
<i>Pachygrapsus transverses</i>	52	6.1	11	1.0	57	3.6	51	1.8	4	0.1	*	*
<i>Callinectes amnicola</i>	3	0.4	41	3.7	5	0.3	52	1.8	9	0.3	*	*
<i>Macrobrachium macrobrachion</i>	11	1.3	15	1.3	*	*	2	0.1	*	*	*	*
Trichoptera												
<i>Phryganea</i> sp. (Caddis fly larva)	1	0.1	1	0.1	1	0.1	1	0.1	*	*	*	*
S =	15	-	16	-	17	-	18	-	16	-	11	-
N =	850	-	1,120	-	1,594	-	2,886	-	3,277	-	1,072	-

S: Number of Species; N: Total number of individuals *: Benthic macroinvertebrates were not encountered

Table 2: Diversity of macrobenthos in the sampling stations of Majidun river

Factors	Station 1 (source of river)	Station 2	Station 3	Station 4	Station 5	Station 6 (mouth of river)
No. of individuals (N)	850	1120	1594	2886	3277	1072
No. of Species (S)	15	16	17	18	16	11
Shannon-wiener diversity (H)	0.880	0.839	0.825	0.971	0.923	0.866
Evenness index (E)	0.748	0.696	0.670	0.773	0.766	0.831

20.2%, closely followed by *Pachymelania aurita* with 19.6% and the least, *Phryganea* sp. a trichoptera insect accounted for <0.1% of the total number of macrobenthos (Fig. 2). Highest gastropods numbers were recorded in August (84.8%) 2008 while the lowest were recorded in July (58.8%) 2008. Trichoptera was only recorded in July and August 2008 and the abundance was <1% of the total number. Bivalvia and Decapoda were relatively constant throughout the study period as shown in the (Fig. 3). The Shannon-wiener diversity index (H) showed that the highest mean of 0.97 was recorded in station 4, followed by station 5 with mean of 0.92 while the lowest mean value (0.83) was recorded in station 3. Evenness index (E) ranged from 0.670 in station 3-0.831 in station 6. The highest number of species (18) was recorded in station 4 while station 6 had the least number (11) species (Table 2).

DISCUSSION

The total number of 18 taxa reported in the present study is far less than those reported for rivers in humid tropics (Bishop, 1973; Edema *et al.*, 2002; Adakole and Anunne, 2003) and these may be as a result of different environmental conditions such as water quality and movement, substrate instability, salinity regime and food

availability just to mention a few (Dance and Hynes, 1980). However, the taxa richness of Majidun river is relatively higher than those of the research of Chukwu and Nwankwo (2003) who recorded 8 taxa in Port Novo Creek and Sikoki and Zabbey (2006) who recorded 14 taxa in Imo river. Gastropods and bivalves are relatively tolerant of physical and chemical variations in the environment and are usually present in a broad range of habitats. It is therefore, not surprising that they dominated the macrobenthos of the study area. Similar research by Ajao and Fagade (2002) also recorded gastropods as the dominant benthic fauna in Lagos Lagoon. The muddy nature of the substratum in the stations may have favours the growths of gastropods (Oyenakan, 1979). Egonmwan (2008) also reported gastropod to inhabit quite water where the substratum is muddy and rich in detritus.

The low abundance of the fresh water prawns, *Macrobrachium macrobrachion* and trichoptera (caddis fly larva) maybe an indication of relatively stressed environmental condition. Abowei *et al.* (2006) reported *Macrobrachium* prawns to be sensitive to deteriorated water quality condition while Emere and Nasiru (2007) reported trichopterans as pollution sensitive macroinvertebrates.

The high Shannon-wiener diversity and evenness value in station 4 indicates high species richness. The low Shannon-wiener value with low evenness in station 3 indicates low species diversity. According to Mackie (1998), the longer a locality has been in the same condition the richer is its biotic community and the more stable it is. Therefore, low diversity in station 3 is an indication of stress in the environment as a result of various human activities while high diversity in station 4 is a reflection of stress-free and stable environment. Similar observations were made by Ajao and Fagade (1990) as to the imbalances of macroinvertebrate from the Western industrialized parts of Lagos Lagoon which received a complex mixture of domestic and industrial waste. The increase in macrobenthos abundance in the months of August to October could be attributed to the life history and population dynamics of the gastropods species that constituted the dominant macrobenthos taxa in this study. They are noted to be adapted to unstable water bodies and exploit brief period of favorable conditions, lasting only a few months to reproduce rapidly and build up dense population (Woolhouse and Chandiwana, 1990). The diversity and abundance of macrobenthos in Majidun river is low. Odum (1971) reported that diversity only tend to be low in physically disturbed ecosystem.

CONCLUSION

The intensity of sand mining and the uncontrolled discharged of domestic and industrial waste have pollution implications as they alter bottom conditions and destroy molluscs settlement areas (Cole, 1977). These may have probably resulted in the low diversity of this tropical river particularly in the 1st 3 stations where many activities occur.

REFERENCES

- Abowei, J.F., S.N. Deekae, M.E. Alison, C.C. Tawari and S.A. Ngodigha, 2006. A Review of Shrimp Fisheries in Nigeria. Pre-Joe Publisher, Port Harcourt, Nigeria, pp: 10.
- Adakole, J.A. and P.A. Anunne, 2003. Benthic macroinvertebrates as indicators of environmental quality of an urban stream in Zaria, Northern Nigeria. J. Aqua. Sci., 18: 85-92.
- Ajao, E.A. and S.O. Fagade, 1990. A study of sediment communities in Lagos Lagoon, Nigeria. J. Oil Chem. Pollut., 7: 85-117.
- Ajao, E.A. and S.O. Fagade, 2002. The benthic Macroinfauna of Lagos Lagoon. Zoologist, 1: 1-5.
- Akpata, T.V.I., J.A. Oyenekan and D.I. Nwankwo, 1993. Impacts of organic pollution on the bacterial, plankton and benthic populations of Lagos Lagoon, Nigeria. Int. J. Ecol. Environ. Sci., 19: 73-82.
- Awosika, L.F. and C.O. Dublin-Green, 1994. Sand mining in the Lagos and Lekki lagoons and strategies for effective management. J. Min. Geol., 30: 137-139.
- Bishop, J.E., 1973. Limnology of a Small Malayan River Sungai Gombak. Dr. W. Junk, Hague, PP: 485.
- Brown, C.A. and E.A. Ajao, 2004. Effects of topographic modification on the composition and abundance of macrofauna in Southern Lagos lagoon. West Afr. J. Applied Ecol., 5: 41-50.
- Celik, K., 2002. Community structure of macrobenthos of a southeast texa sand-pit lake related to water temperature, pH and Dissolved oxygen concentration. Turk. J. Zool., 26: 333-339.
- Chukwu, L.O. and D.I. Nwankwo, 2003. The impact of land based pollution on the hydrochemistry and macrobenthic community of a tropical West African creek. Proceeding of 7th International Specialised Conferene on Diffuse Pollution and Basin Manag, 2003, Dublin, Ireland, pp: 1-16.
- Cole, H.A., 1977. Sand and gravel extraction. Marine Polut. Bull., 8: 193-240.
- Dance, K.W. and H.B.N. Hynes, 1980. Some effects of agricultural land use on stream insect communities. Environ. Pollut. Ser. A. Ecol. Biol., 22: 19-28.
- Diaz, R.J., M. Solan and R.M. Valente, 2004. A review of approaches for classifying benthic habitats and evaluating habitat quality. J. Environ. Manage., 73: 165-181.
- Dillon Jr, R.T., 2000. The Ecology of Feshwater Molluscs. Cambridge University Press, Cambridge, UK.
- Edema, C.U., J.O. Ayeni and A. Aruoture, 2002. Some observations on the zooplankton and macrobenthos of the Okhuo River, Nigeria. J. Aqua. Sci., 17: 145-149.
- Edmunds, J., 1978. Sea Shells and other Molluscs found on West African Coast and Estuaries. Ghana University Press, Accra, pp: 146.
- Edokpayi, C.A., J.C. Okenyi, A.E. Ogbeibu and E.C. Osimen, 2000. The effect of human activities on the macrobenthic invertebrates of Ibiekuma Stream, Ekpoma, Nigeria. Biosci. Res. Commun., 12: 79-87.
- Egborge, A.B.M., L.I. Ezemonye and G.E. Awoze, 2003. Macroinvertebrate fauna of Udu-Ughievwen Wetlands, Southern, Nigeria. J. Aqua. Sci., 18: 1-8.
- Egonmwan, R.I., 2008. The ecology and habits of *Tympanotonus fuscatus* var. *radula* (L.) (Prosobranchia: Potamididae). J. Boil. Sci., 8: 186-190.
- Emere, M.C. and C.E. Nasiru, 2007. Macroinvertebrate as indicators of the water quality of an urbanised Stream, Kaduna, Nigeria. Medwell J. Fish. Int., 2: 152-157.

- George, A.D.I., J.F.N. Abowei and E.R. Daka, 2009. Benthic macroinvertebrate fauna and physico-chemical parameters in Okpoka creek sediments, Niger Delta, Nigeria. *Int. J. Anim. Vet. Adv.*, 1: 59-65.
- Holme, N.A. and A.D. McIntyre, 1984. *Methods for the Study of Marine Benthos*. 2nd Edn., Blackwell Scientific Publications, Oxford, London, Edinburgh, pp: 387.
- Ikomi, R.B., F.O. Arimoro and O.K. Odihirin, 2005. Composition, distribution and abundance of macroinvertebrates of the upper reaches of River Ethiope, Delta State, Nigeria. *The Zoologist*, 3: 68-81.
- Lloyd, M. and R.G. Ghellardi, 1964. A table for calculating the equitability components of species diversity. *J. Anim. Ecol.*, 33: 217-225.
- Lydeard, C., R.H. Cowie, W.F. Ponder, A.E. Bogan and P. Bouchet, 2004. The global decline of nonmarine mollusks. *Bioscience*, 54: 321-330.
- Macan, T.T., 1959. *A Guide to Freshwater Invertebrate Animals*. Longman Group Ltd., England.
- Mackie, G.L., 1998. *Applied Aquatic Ecosystem Concepts*. University of Guelph Custom Coursepack, 12 Chapters, Index, Canada.
- Odum, E.P., 1971. *Fundamentals of Ecology*. 3rd Edn., W.B. Saunders Company, Philadelphia, pp: 574.
- Oyenakan, J.A., 1979. The ecology of the genus *Pachymelania* in Lagos lagoon. *Arch. Hydrobiol.*, 86: 515-522.
- Pennak, R.W., 1978. *Freshwater Invertebrates of the United States*. 2nd Edn., John Wiley and Sons, USA., pp: 803.
- Rosenberg, D.M. and V.H. Resh, 1993. *Freshwater Biomonitoring and Benthic Invertebrate*. Chapman and Hall, New York.
- Shannon, C.E. and W. Weaver, 1963. *The Mathematical Theory of Communications*. University of Illinois Press, Urbana, pp: 125.
- Sikoki, F.D. and N. Zabbey, 2006. Environmental gradients and Benthic community of the middle reaches of Imo River, South-Eastern Nigeria. *Environ. Ecol.*, 24: 32-36.
- WHO, 1978. *A Field Guide to African Freshwater Snails*. WHO Snail Identification Center, Danish Bilharziasis Lab. Jaegersborg, Charlottenlund Denmark, pp: 30.
- Woolhouse, M.E. and S.K. Chandiwana, 1990. Population dynamics model for *Bulinus globosus*, intermediate host for *Schistosoma haematobium*, in river habitats. *Acta Trop.*, 47: 151-160.